

Impact of exercise endurance training on *purβ* gene expression and cardiac function

Maryam Anjomshoa¹, Naser Rostamzadeh², Omid Rostamzadeh³, Ayoob Rostamzadeh^{1*}

¹Department of Anatomical Sciences, Shahrekord University of Medical Sciences, Shahrekord, Iran.

²Department of Physical Education and Sport Sciences, Shahid Rajaei Teacher Training University, Tehran, Iran.

³Department of Occupational Therapy, Faculty of Rehabilitation Sciences, Iran University of Medical Sciences, Tehran, Iran.

*Corresponding author: Ayoob Rostamzadeh, Department of Anatomical Sciences, Faculty of Medicine, Shahrekord University of Medical Sciences, Rahmatiyeh Town, Shahrekord, Iran. Tel: +989187225635. Email: ayoobrostamzade@gmail.com, Fax: +983813334911.

Abstract

Introduction: Endurance training has significant effects on the renewal of heart tissue, including myosin heavy chain (MHC) proteins. On the other side, Purine-rich element-binding protein β (*purβ*) decreases the α MHC gene expression. The aim of this study was to determine the impact of exercise endurance training on *purβ* gene expression in the heart of Wistar rats.

Methods: Fourteen rats have been kept under controlled conditions and after familiarizing with training protocol, they were divided into control groups and experimental groups. The experimental group performed a 10-week treadmill running program for 30 min/day, 5 days/week. 48 hours after the last training session, the rats were anesthetized and the heart and their left ventricle were taken out and *purβ* expression was measured using real time PCR method. All data were analyzed using *t* test.

Results: In this study, the results of M-mode echocardiography showed that endurance training led to cardiac hypertrophy. After endurance training, the heart weight, especially the left ventricular weight significantly increased. The *purβ* gene expression significantly decreased in the left ventricular tissue of endurance-trained rats.

Conclusion: The results of this study revealed that endurance training has considerable effects on heart size and *purβ* gene expression. The *purβ* gene also repressed α MHC gene expression; it seems that the changes in heart structure related to α MHC gene expression.

Keywords: Gene expression, *purβ* gene, Cardiac plasticity, Endurance training

Introduction

In response to different environmental demands, cardiac structures and functions considerably changed. Physical activities have a drastic effect on heart remodeling and gene expression in heart tissue (1). Heart adaptation in response to endurance training is related to the variety of stimuli imposed to the heart, so morphological changes in the heart are different in response to each activity (2). Adaptation of athlete's heart to endurance training includes changes such as increased wall thickness and left ventricular mass (3, 4). Also relative cavity diameter of heart in endurance runners is higher compared to sprint runners (3). Structural changes in left ventricle of heart demonstrated adaptation to hemodynamic overload induced by endurance training. Work capacity during exercise is positively influenced by preload increase after endurance training, while increased afterload due to isometric training in strength-trained athletes, determines higher systemic resistance during physical effort (4). There are different signaling pathways in heart that

mediate the heart growth response. These signaling circuits directly control hypertrophic growth by altering gene expression in cardiac myocytes (5). There are wide varieties of physiological and pathological stimuli that affect the relative proportions of the two forms of the motor protein myosin heavy chain (MHC). The expression of faster MHC motor protein, α , (α MHC) in heart leads to produce more power than slower MHC motor protein, β , (β MHC) and so the heart power and contractility would increase (6). In left ventricle of normal hearts, the α MHC mRNA was expressed at considerable levels and in the end stage of heart failure, the α MHC mRNA expression, decreases up to 15-fold and also these changes would occur in the level of MHC protein. In nonfailing hearts, the α MHC protein represented about 7% of the total MHC protein, while in failing hearts, there was effectively no detectable α MHC protein in the left ventricles (6). No myocardial collagen produces in hypertrophy caused by exercise (7). Physiological and pathological cardiac hypertrophy has opposite changes in β MHC gene expressions (8).

The Purine-rich element-binding (*purβ*) protein is single-strand DNA binding proteins that can bind to DNA as the homodimer or heterodimer. This protein participates in cell differentiation, proliferation and apoptosis and cell-specific gene regulation (9). The *purβ* mediate repression of β MHC gene expression. The levels of *purβ* in heart failure would increase and is important in regulating the transcription of α MHC gene (10). The aim of this study was to investigate the effects of 10 weeks of endurance training on *Purβ* gene expression in the heart tissue of Wistar rats.

Material and methods

Animals care

Fourteen male Wistar rats of 5 weeks old (200 ± 20 gr) were obtained from the Pasteur institute (Tehran, Iran). All animals were kept at room temperature ($22 \pm 3^\circ\text{C}$) with a 12-h light/dark cycle until they reach puberty. Within this time, rats were maintained in 4 equal cages. At the end of this step, the average and standard deviation of rat's weights was 231 ± 24 gr. All rats were familiarized with treadmill running for 5 sessions/10 days. At the end of this session, rats were divided randomly into two groups, experimental group and control group (7 rats in each group).

Training protocol

According to previous studies, we designed an endurance training protocol for rats as previously described (11, 12), so that endurance training could cause cardiac hypertrophy. The rats trained 6 days/week for 4 weeks with speed, grade and duration progressively increased. To prevent the rats from stopping, an electric grid at the rear of the belt was used as the running stimulus. The rats began training at 12 m/min and 0% grade for 5 min in first day. By the 2nd weeks duration and speed were increased until the rats ran at 30 m/min for 60 min/day. During 5–8 weeks the speed and duration were maintained and grade was gradually increased to 5%. Finally, 48 hours after the end of the last exercise session, rats were anesthetized with a combination of Ketamine (50 mg/kg) and Xylazine (5 mg/kg) and under sterile surgical methods; their heart and left ventricle were isolated and placed into 1.5 ml microtubes. Then heart tissue explants were moved to a liquid nitrogen tank for

cryopreservation. The frozen heart explants were homogenized in liquid nitrogen using a mortar. In this study, m-mode was used to demonstrate the effects of endurance training on the heart structure. Furthermore, many studies used heart weight/body weight ratio, and heart weight/body surface area (BSA) ratio to evaluate cardiac hypertrophy (13, 14). In this study to evaluate the cardiac hypertrophy, two factors were used for normalization the left ventricular weight. Weight and length of animal (from mouth to root of tail) were measured in the anesthetized state for BSA calculations (15). The heart and left ventricle were weighed separately using a digital analytical balance (Sartorius, model-BL210S; Japan). BSA of rats was evaluated using the following formula:

$$\text{BSA} = 6.67 \times W^{0.7} \times [0.34 / (\sqrt[3]{W/L})]. \quad W = \text{body weight (gr)}; L = \text{body length (cm)}$$

Animals and ethics

Animal experimentations were approved by the Ethical Committee of Shahid Rajaei Teacher training University and carried out in an ethically proper way by following the provided guidelines.

RNA extraction

Total RNA was extracted from heart tissue as previously described (16). Briefly, 1 ml TRIZOL Reagent (Invitrogen, Carlsbad, CA, USA) was added to 100 mg heart tissue. Then the RNA samples were treated with RNase free DNase to remove any residual DNA.

First-strand cDNA synthesis and Real-Time Quantification of Gene Expression

Total extracted RNA was transcribed to the first strand complementary DNA (cDNA) with the cDNA Synthesis Kit (Thermo Scientific, Schwerte, Germany) according to manufacturer's instructions. Real time RT-PCR was performed using the Rotor Gene System (Applied Biosystems, Darmstadt, Germany).

First-strand cDNA synthesis and Real-Time Quantification of Gene Expression

Total extracted RNA was transcribed to the first strand complementary DNA (cDNA) with the cDNA Synthesis Kit (Thermo Scientific, Schwerte, Germany) according to manufacturer's instructions. Real time RT-PCR was performed using the Rotor Gene System (Applied Biosystems, Darmstadt, Germany). Each reaction mix contained 1 μ l cDNA, 10 μ l SYBR green master mix (TaKaRa, Kyoto, Japan) and 5 μ l forward and reverse primers (Table 1) adjusted to a total volume of 20 ml using distilled water. The transcripts of *pur β* along with Glyceraldehyde 3-Phosphate Dehydrogenase (*gapdh*), as a reference gene (17), were analyzed by quantitative real time RT-PCR. The primers characteristics are shown in Table 1. For each cDNA sample, three replicates of PCR were carried out. CT sample of *pur β* gene were normalized to the CT of *gapdh* represented as $2^{-\Delta\Delta CT}$ (18-20).

Name		Sequence 5-3	NCBI Reference Sequence	Product size
<i>Gap dh</i>	F	AACCCATCACCATCTTCCAG	NM_017008.4	74
	R	CACGACATACTCAGACCAG		
<i>Purβ</i>	F	GTGAGGAAGTGGATGAGGATTG	NM_001017503.1	100
	R	GGACGAGTAGGAAAGGGAAC		

Statistical analysis

The transformed data of RT-PCR expression of *pur β* gene were analyzed by Shapiro-Wilk test of SPSS 20. Then the differences between groups were compared by *t* test. Differences were considered as significant at $P < 0.05$.

Results

The results of this study showed that after 10 weeks of endurance training, a hypertrophy occurred in left ventricle, which was supported by evaluating the left ventricular weight/body weight ratio and left ventricular weight/body surface ratio (Figures 1 and 2). Figure 1 show the left ventricular weight/BSA ratio in the experimental group (0.168 ± 0.008) is significantly higher in comparison with control group (0.153 ± 0.006) ($p = 0.006$). On the other hand, figure 2 revealed that the heart and left ventricular weight of experimental group is higher than the control group,

so that the left ventricular weight/ body weight ratio in experimental group was significantly higher ($2/3 \pm 0.18$) compared to control group (2.049 ± 0.12) ($p = 0.04$).




Figure 1. Analysis of hypertrophy indexes show that the left ventricular weight/ BSA ratio in the experimental group, is higher compared to control group ($P = 0.006$). Error bars shows 95% CI.




Figure 2. Analysis of hypertrophy indexes show that the left ventricular weight/ body weight ratio in experimental group, is higher compared to control group ($P = 0.04$). Error bars shows 95% CI.

Also, the results of the *t*-test ($t = -4.35$) demonstrated that the average *Pur β* gene expression in heart of experimental group significantly decreased to 54% in comparison with control group after 10 weeks of endurance training ($p = 0.008$) (Figure 3).




Figure 3. Results demonstrated that the average *Pur β* gene expression in heart of experimental group decreased in contrast to control group after 10 weeks of endurance training ($P = 0.008$). Error bars shows 95% CI.

Discussion

Currently, the various aspects of exercise-induced cardiac remodeling, including ventricular chamber enlargement, myocardial hypertrophy have been recognized (21), but the cellular pathways responsible for cardiac remodeling remain poorly understood. The results of the present study demonstrated that endurance training led to reduction of *Purβ* gene expression in left ventricle to 46%. To date, except this study any previous information has been reported about a role endurance training on *purβ* gene expression, so the results of the present study was discussed based on the studies that identified other effects of this gene. In the rat, α MHC is the predominant isoform in adult hearts, has high ATPase activity, and is associated with increased shortening velocity of the cardiac myocytes. Therefore, changes in the α MHC in the cardiac ventricle alter the contractile properties of the heart (22). Miyata et al. (2000) showed that α MHC mRNA was expressed at considerable levels in the nonfailing heart and was considerably decreased in heart failure and pathological cardiac hypertrophy (6). van Rooij et al. (2007) showed that down-regulation of α MHC gene is a process that occurs in the pathological cardiac hypertrophy (23). The *purβ* protein is a single-strand DNA binding protein that can bind to DNA as the homodimer or heterodimer. The *purβ* protein is also highly expressed in cardiac myocytes, and their level changes in heart failure. Gupta et al. (2003) described Overexpression of *purβ* down regulates α MHC gene expression in heart tissue (9). The *purβ* protein cooperated with other cardiac factors binding to the α MHC gene promoter in a negative manner. The *purβ* protein has been shown to bind to other cardiac myogenic factors which also play prominent roles in α MHC gene expression. The studies of Gupta et al. (2003) also showed that *purβ* protein is capable of binding to α MHC mRNA and attenuate its translational efficiency and expression of *purβ* protein in failing hearts suppressed the α MHC mRNA (9). In cell renewal, the activity of *Purβ* gene and the level of protein increase, however, it decreases in response to mechanical overload. In heart failure the *Purβ* level increased, which is contributed to the transcription of α MHC (10). The results of McCarthy's study confirm that increase of *Purβ* expression led to muscles atrophy (24). Stevenson et al. (2003) showed that suppression of MHC genes associated with increased expression of *Purβ* gene

(25). Comparing the results of McCarthy with the present report demonstrated that *Purβ* has major roles in the cardiac myocytes in response to cardiac hypertrophy. In the present study, we did not evaluate the *Purβ* protein level that is the final factor for the effect of endurance training on the heart tissue, so for future studies, evaluating the *Purβ* protein level in heart tissue after endurance training would be recommended. In conclusion, the results of the present study demonstrated that endurance training decreased the *Purβ* gene expression. Cardiac remodeling (including changes in dimensions, mass and shape of the hearts) in response to endurance training is in association with neurohormonal activation and changes in of heart size, specially left ventricle, which in turn. Considering different effects of endurance training on the cardiac hypertrophy, *Purβ* gene expression and also the effect of *Purβ* in the α MHC expression, it seems that heart remodeling in response to endurance training is related to α MHC expression.

Acknowledgements

The authors would like to thank the staff of the deputy research in the Shahid Rajaei Teacher training University for their kind cooperation.

Conflict of Interest

The authors declare that they have no conflict of interests.

References

- Hill JA, Olson EN. Cardiac plasticity. *N Engl J Med*. 2008;358(13):1370-80.
- Pluim BM, Zwinderman AH, van der Laarse A, van der Wall EE. The athlete's heart: a meta-analysis of cardiac structure and function. *Circulation*. 2000;101(3):336-44.
- Venckunas T, Raugaliene R, Mazutaitiene B, Ramoskeviciute S. Endurance rather than sprint running training increases left ventricular wall thickness in female athletes. *Eur J Appl Physiol*. 2008; 102(3):307-11.
- D'Andrea A, Limongelli G, Caso P, Sarubbi B, Della Pietra A, Brancaccio P, et al. Association between left ventricular structure and cardiac performance during effort in two morphological forms of athlete's heart. *Int J Cardiol*. 2002;86(2-3):177-84.
- Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signalling pathways. *Nat Rev Mol Cell Biol*. 2006;7(8):589-600.
- Miyata S, Minobe W, Bristow MR, Leinwand LA. Myosin heavy chain isoform expression in the failing and nonfailing human heart. *Circ Res*. 2000;86(4):386-90.

7. Dorn GW, 2nd. The fuzzy logic of physiological cardiac hypertrophy. *Hypertension*. 2007;49(5):962-70.
8. Kinugawa K, Yonekura K, Ribeiro RC, Eto Y, Aoyagi T, Baxter JD, et al. Regulation of thyroid hormone receptor isoforms in physiological and pathological cardiac hypertrophy. *Circ Res*. 2001;89(7):591-8.
9. Gupta M, Sueblinvong V, Raman J, Jeevanandam V, Gupta MP. Single-stranded DNA-binding proteins PURalpha and PURbeta bind to a purine-rich negative regulatory element of the alpha-myosin heavy chain gene and control transcriptional and translational regulation of the gene expression. Implications in the repression of alpha-myosin heavy chain during heart failure. *J Biol Chem*. 2003;278(45):44935-48.
10. Ji J, Tsika GL, Rindt H, Schreiber KL, McCarthy JJ, Kelm RJ Jr., et al. Puralpha and Purbeta collaborate with Sp3 to negatively regulate beta-myosin heavy chain gene expression during skeletal muscle inactivity. *Mol Cell Biol*. 2007;27(4):1531-43.
11. Jin H, Yang R, Li W, Lu H, Ryan AM, Ogasawara AK, et al. Effects of exercise training on cardiac function, gene expression, and apoptosis in rats. *Am J Physiol Heart Circ Physiol*. 2000;279(6):2994-3002.
12. Sun L, Shen W, Liu Z, Guan S, Liu J, Ding S. Endurance exercise causes mitochondrial and oxidative stress in rat liver: effects of a combination of mitochondrial targeting nutrients. *Life Sci*. 2010;86(1-2):39-44.
13. Zhu SS, Ma JZ, Yong YH, Niu J, Zhang JN. Left ventricular function in physiologic and pathologic hypertrophy in Sprague-Dawley rats. *Science & Sports*. 2008; 23(6): 299-305.
14. Seo JS, Lee SY, Won KJ, Kim DJ, Sohn DS, Yang KM, et al. Relationship between normal heart size and body indices in Korean. *J Korean Med Sci*. 2000;15(6):641-6.
15. Haddad R, Kasneci A, Mephram K, Sebag IA, Chalifour LE. Gestational exposure to diethylstilbestrol alters cardiac structure/function, protein expression and DNA methylation in adult male mice progeny. *Toxicol Appl Pharmacol*. 2013; 266(1): 27-37.
16. Ghanbari-Niaki A, Ghanbari-Abarghoi S, Rahbarizadeh F, Zare-Kookandeh N, Gholizadeh M, Roudbari F, et al. Heart ABCA1 and PPAR- α Genes Expression Responses in Male rats: Effects of High Intensity Treadmill Running Training and Aqueous Extraction of Black Crataegus-Pentaegyna. *Res Cardiovasc Med*. 2013; 2(4): 153-159.
17. Silver N, Cotroneo E, Proctor G, Osailan S, Paterson KL, Carpenter GH. Selection of housekeeping genes for gene expression studies in the adult rat submandibular gland under normal, inflamed, atrophic and regenerative states. *BMC Mol Biol*. 2008; 9:64.
18. Yuan JS, Reed A, Chen F, Stewart CN Jr. Statistical analysis of real-time PCR data. *BMC Bioinformatics*. 2006; 7:85-97.
19. Wong ML, Medrano JF. Real-time PCR for mRNA quantitation. *Biotechniques*. 2005; 39(1):75-85.
20. Gunning P, O'neill G, Hardeman E. Tropomyosin-based regulation of the actin cytoskeleton in time and space. *Physiol Rev*. 2008; 88(1):1-35.
21. Weiner RB, Baggish AL. Exercise-induced cardiac remodeling. *Prog Cardiovasc Dis*. 2012; 54(5):380-6.
22. Kim S, Iwao H. Molecular and Cellular Mechanisms of Angiotensin II-Mediated Cardiovascular and Renal Diseases. *Pharmacol Rev*. 2000; 52(1): 11-34.
23. van Rooij E, Sutherland LB, Qi X, Richardson JA, Hill J, Olson EN. Control of stress-dependent cardiac growth and gene expression by a microRNA. *Science*. 2007;316(5824):575-9.
24. McCarthy JJ, Esser KA, Peterson CA, Dupont-Versteegden EE. Evidence of MyomiR network regulation of beta-myosin heavy chain gene expression during skeletal muscle atrophy. *Physiol Genomics*. 2009;39(3):219-26.
25. Stevenson EJ, Giresi PG, Koncarevic A, Kandarian SC. Global analysis of gene expression patterns during disuse atrophy in rat skeletal muscle. *J Physiol*. 2003;551(Pt 1):33-48.